

THE ACTIVE PHARM · EST. 2024



The Science of Human Potential

SINGLE PEPTIDE CATALOG

VOLUME II · SINGLE AGENTS

TWENTY-FIVE COMPOUNDS

Contents

Twenty-Five single agents, organized by pharmacological function. Each title links to its full profile.

METABOLIC & FAT LOSS

01	Semaglutide	GLP-1 Receptor Agonist	p. 3
02	Tirzepatide		p. 4
	GIP / GLP-1 Receptor Agonist		
03	AOD-9604	HGH Fragment Analog	p. 5
04	5-Amino-1MQ	NNMT Inhibitor	p. 6

GROWTH HORMONE & ANABOLIC SIGNALING

05	Sermorelin	GHRH Analog	p. 7
06	Tesamorelin	Stabilized GHRH Analog	p. 8
07	Ipamorelin		p. 9
	Growth Hormone Secretagogue		
08	IGF-1 LR3	Insulin-like Growth Factor	p. 10

MITOCHONDRIAL & CELLULAR ENERGY

09	MOIS-C	Mitochondrial Peptide	p. 11
10	NAD+	Cellular Coenzyme	p. 12
11	SS-31	Mitochondrial-Targeting Peptide	p. 13
12	Glutathione		p. 14
	Master Antioxidant Tripeptide		

TISSUE REPAIR & RECOVERY

13	BPC-157	Body Protection Compound	p. 15
14	TB-500	Thymosin Beta-4 Fragment	p. 16
15	GHK-Cu	Copper Tripeptide	p. 17

INFLAMMATION & IMMUNE DEFENSE

16	Thymosin Alpha-1		p. 18
	Immune Modulator		
17	KPV	Anti-inflammatory Tripeptide	p. 19
18	LL-37	Antimicrobial Peptide	p. 20

COGNITIVE FUNCTION & NEUROPROTECTION

19	Semax	Nootropic Peptide	p. 21
20	Selank	Anxiolytic Peptide	p. 22

LONGEVITY, SLEEP & CIRCADIAN

21	Epitalon	Telomerase Activator	p. 23
22	DSIP	Delta Sleep-Inducing Peptide	p. 24

SEXUAL & REPRODUCTIVE HEALTH

23	PT-141	Melanocortin Agonist	p. 25
24	Oxytocin	Neuropeptide Hormone	p. 26
25	Kisspeptin	Reproductive Signal	p. 27



Semaglutide

The incretin signal for weight and glucose control.

GLP-1 RECEPTOR AGONIST

01



HOW IT WORKS

Activates GLP-1 receptors to enhance glucose-dependent insulin secretion, suppress glucagon, and slow gastric emptying, while signaling satiety in the hypothalamus to reduce appetite and caloric intake.

02



WEIGHT LOSS

Produces substantial, sustained weight loss in clinical trials.

03



GLYCEMIC CONTROL

Lowers blood glucose and HbA1c with low hypoglycemia risk.

MOLECULAR FORMULA

GLP-1

receptor agonist

$C_{187}H_{291}N_{45}O_{59}$

MW $\approx$ 4,113.6 Da

5 MG / 2 ML VIAL

GLP-1 Receptor Agonist

04



PAIRS WELL WITH

Ipamorelin · preserves lean mass during weight loss

Tesamorelin · visceral fat and lean-mass support

AOD-9604 · enhanced fat metabolism


Complementary agents commonly combined in clinical protocols.

05



CARDIOMETABOLIC

Associated with improved cardiovascular and metabolic markers.



METABOLIC & FAT LOSS



SAFETY NOTES

Generally well tolerated. The most common effects are gastrointestinal: nausea, vomiting, diarrhea, or constipation, usually transient and tending to ease over time. Less common concerns include gallbladder issues and pancreatitis. Contraindicated with a personal or family history of medullary thyroid carcinoma or MEN2. Because rapid weight loss can include lean mass, resistance training and muscle-preserving agents are commonly paired.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

Tirzepatide

Two incretin pathways, activated at once.

DUAL GIP AND GLP-1 RECEPTOR AGONIST

01



HOW IT WORKS

Co-activates GIP and GLP-1 receptors to amplify insulin release, suppress glucagon, slow gastric emptying, and reduce appetite, enhancing weight loss and glycemic control beyond GLP-1 alone.

02



WEIGHT LOSS

Among the largest weight reductions reported in incretin trials.

03



GLYCEMIC CONTROL

Strong reductions in blood glucose and HbA1c.

MOLECULAR FORMULA

GLP-1·GIP

dual receptor agonist

$C_{22}H_{34}N_4O_6$

MW ≈ 4,813.5 Da

10 MG / 2 ML VIAL

Dual GIP and GLP-1 Receptor
Agonist

04



PAIRS WELL WITH

Ipamorelin · preserves lean mass during weight loss
Tesamorelin · visceral fat and lean-mass support
AOD-9604 · enhanced fat metabolism


Complementary agents commonly combined in clinical protocols.

05



DUAL INCRETIN

Engages both GIP and GLP-1 pathways for an amplified effect.



METABOLIC & FAT LOSS



SAFETY NOTES

Generally well tolerated. Gastrointestinal effects (nausea, diarrhea, constipation) are most common and tend to ease over time. Less common concerns include gallbladder issues and pancreatitis. Contraindicated with a personal or family history of medullary thyroid carcinoma or MEN2. As with any rapid weight loss, lean-mass loss is a concern.



AOD-9604

The fat-loss fragment of growth hormone.

MODIFIED HGH 176-191 FRAGMENT

01



HOW IT WORKS

Stimulates beta-adrenergic receptors on fat cells to release stored triglycerides while blocking new fat formation, without touching the growth hormone or insulin pathways.

02



FAT REDUCTION

Targets abdominal and visceral fat for reduction.


03



ANTI-LIPOGENIC

Inhibits new fat formation without hormonal load.

MOLECULAR STRUCTURE



C₇₈H₁₂₅N₂₃O₂₃S₂

MW ≈ 1,817.1 Da

10 MG / 5 ML VIAL

Modified HGH 176-191
Fragment

04



PAIRS WELL WITH

GLP therapy · complements incretin-driven fat loss
5-Amino-1MQ · energy expenditure
MOTS-c · metabolic


Complementary agents commonly combined in clinical protocols.

05



JOINT SUPPORT

Offers joint and anti-inflammatory support.



METABOLIC & FAT LOSS



SAFETY NOTES

A high safety profile in clinical trials: non-toxic, non-immunogenic, and non-hormonal, with no significant reported side effects and no suppression of natural growth hormone or insulin function. Effects appear only with consistent use over two to three months; a single month is generally insufficient for noticeable fat loss.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

5-Amino-1MQ

Releasing the brake on metabolism.

NNMT ENZYME INHIBITOR

01



HOW IT WORKS


Inhibits the enzyme NNMT to regulate cellular NAD+ metabolism, shifting signaling toward greater energy expenditure and metabolic efficiency in fat tissue rather than altering hormones or appetite.

02



METABOLIC EFFICIENCY

Supports healthy metabolism, energy use, and reduced fat accumulation.



MW $\approx$ 286.1 Da

10 MG / 5 ML VIAL

NNMT Enzyme Inhibitor

03



PAIRS WELL WITH

AOD-9604 · fat metabolism
MOTS-c · mitochondrial
NAD+ · cellular energy

Complementary agents commonly combined in clinical protocols.

04



ENZYMATIC PATHWAY

Acts on the NNMT enzyme rather than on hormones.



METABOLIC & FAT LOSS



SAFETY NOTES

As an NNMT inhibitor used in experimental metabolic protocols, 5-Amino-1MQ is less standardized than many peptides. Providers should watch for stimulation-type effects such as restlessness or insomnia, GI upset, headache, or appetite changes. Use caution in patients with significant metabolic disease until tolerance and monitoring are established.



Sermorelin

THE ACTIVE PHARM

EST. 2024

503A PHARMACY COMPOUNDED

Growth hormone, prompted not replaced.

GROWTH HORMONE-RELEASING HORMONE ANALOG

01



HOW IT WORKS

Binds pituitary GHRH receptors to release the body's own growth hormone and downstream IGF-1, working with the natural nighttime rhythm and preserving normal pituitary feedback.

02



GH OUTPUT

Raises natural growth hormone output without direct replacement.


03



SLEEP & RECOVERY

Improves sleep depth, recovery, and tissue regeneration.

MOLECULAR STRUCTURE



C₁₄₉H₂₄₆N₄₄O₄₂S

MW ≈ 3,357.9 Da

10 MG / 5 ML VIAL

Growth Hormone-Releasing
Hormone Analog

04



PAIRS WELL WITH

GLP therapy · lean-mass support during weight loss
Ipamorelin · complementary GH pulse
BPC-157 · recovery


Complementary agents commonly combined in clinical protocols.

05



BODY & SKIN

Supports lean mass, body composition, and skin quality.



**GROWTH HORMONE &
ANABOLIC SIGNALING**



SAFETY NOTES

Sermorelin carries a strong safety profile. Side effects are generally mild: transient flushing, dizziness, injection-site irritation, or headache, with occasional increased appetite or vivid dreams from heightened growth hormone output. It is typically administered subcutaneously at night to align with circadian release, and effects accumulate over several weeks as IGF-1 rises gradually.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

Tesamorelin

Targeted reduction of visceral fat.

STABILIZED GROWTH HORMONE-RELEASING HORMONE ANALOG

01



HOW IT WORKS

A stabilized GHRH analog that drives pulsatile growth hormone and IGF-1; a hexenoic-acid group resists breakdown for higher, sustained release and targeted visceral-fat loss.

02



VISCERAL FAT

Significantly reduces visceral adipose tissue.


03



LEAN MASS

Supports lean mass and body recomposition.

MOLECULAR STRUCTURE



C₂₂₀H₃₆₆N₇₂O₆₇S

MW ≈ 5,123.8 Da

10 MG / 5 ML VIAL

Stabilized Growth Hormone-Releasing Hormone Analog

04



PAIRS WELL WITH

GLP therapy · lean-mass support during weight loss

Ipamorelin · GH pulse
MOTS-c · metabolic


Complementary agents commonly combined in clinical protocols.

05



METABOLIC MARKERS

May improve insulin sensitivity and lipid profiles.



**GROWTH HORMONE &
ANABOLIC SIGNALING**



SAFETY NOTES

Well studied with a strong safety profile. Common effects are injection-site irritation, mild edema, muscle stiffness, or joint discomfort tied to elevated growth hormone. It does not suppress natural production and does not cause acromegaly when used appropriately. Use under supervision in patients with diabetes, active malignancy, or cardiovascular disease.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

Ipamorelin

A clean pulse of growth hormone.

SELECTIVE GROWTH HORMONE SECRETAGOGUE

01



HOW IT WORKS

Activates pituitary GHS-R, the ghrelin receptor, for a clean, pulsatile growth hormone release without the broad hormonal disturbance of less selective agents.

02



MUSCLE & RECOVERY

Supports lean muscle and recovery from training or injury.


03



SLEEP & METABOLISM

Improves sleep quality and healthy fat metabolism.

MOLECULAR STRUCTURE



$C_{38}H_{49}N_9O_5$

MW $\approx$ 711.9 Da

10 MG / 5 ML VIAL

Selective Growth Hormone
Secretagogue

04



PAIRS WELL WITH

GLP therapy ·
preserves lean mass
during weight loss
Tesamorelin · visceral
fat
BPC-157 · recovery


Complementary agents
commonly combined in
clinical protocols.

05



SELECTIVE ACTION

Raises GH and IGF-1
signaling while sparing
cortisol and prolactin.



GROWTH HORMONE &
ANABOLIC SIGNALING



SAFETY NOTES

Generally well tolerated. Possible effects include transient flushing, headache, lightheadedness, water retention, and increased appetite. Because it raises growth hormone and IGF-1 signaling indirectly, use clinician oversight in patients with diabetes or insulin resistance, active malignancy, or significant edema or cardiovascular risk.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

IGF-1 LR3

Growth hormone's anabolic message, delivered directly.

INSULIN-LIKE GROWTH FACTOR 1, LONG ARG3

01



HOW IT WORKS

Binds the IGF-1 receptor to activate PI3K/Akt and MAPK signaling for protein synthesis and cell growth; resists binding proteins for a long, sustained anabolic signal.

02



LEAN MUSCLE

Increases lean muscle through enhanced protein synthesis.


03



RECOVERY & REPAIR

Accelerates recovery and supports tendon, ligament, and connective-tissue repair.

MOLECULAR STRUCTURE



C₃₉₀H₆₂₅N₁₁₁O₁₁₅S₅

MW ≈ 8,869.3 Da

10 MG / 5 ML VIAL

Insulin-like Growth Factor 1,
Long Arg3

04



PAIRS WELL WITH

Ipamorelin · GH pulse
BPC-157 · tissue repair
TB-500 · connective tissue

Complementary agents commonly combined in clinical protocols.

05



NUTRIENT UPTAKE

Improves nutrient partitioning and cellular hydration.



**GROWTH HORMONE &
ANABOLIC SIGNALING**



SAFETY NOTES

A highly potent compound that requires careful clinical oversight. Excess can cause hypoglycemia, organ enlargement, or accelerated growth of existing malignancy given its strong proliferative action. It should not be used by anyone with a history of cancer without physician clearance, and blood glucose should be monitored throughout use.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

MOTS-c

A signal written in mitochondrial DNA.

MITOCHONDRIAL-DERIVED PEPTIDE

01



HOW IT WORKS

Encoded by mitochondrial DNA, MOTS-c activates AMPK, the cell's metabolic master switch, promoting fat oxidation, glucose use, and mitochondrial biogenesis, and acting as a mitokine to distant tissues.

02



GLUCOSE CONTROL

Improves insulin sensitivity and glucose control.


03



BODY COMPOSITION

Reduces visceral fat and supports body composition.

MOLECULAR STRUCTURE



C₄₉H₇₅N₁₇O₁₅

MW ≈ 1,142.2 Da

10 MG / 5 ML VIAL

Mitochondrial-Derived
Peptide

04



PAIRS WELL WITH

NAD+ · cellular energy
AOD-9604 · fat metabolism
Tesamorelin · metabolic


Complementary agents commonly combined in clinical protocols.

05



ENDURANCE

Enhances endurance and buffers fatigue.



**MITOCHONDRIAL &
CELLULAR ENERGY**



SAFETY NOTES

A strong safety profile in preclinical and limited human research. Side effects are rare and may include temporary fatigue or mild injection-site irritation. Because it modulates key metabolic pathways, use caution alongside blood-sugar-lowering medications to avoid hypoglycemia.



NAD+

The coenzyme behind cellular energy.

NICOTINAMIDE ADENINE DINUCLEOTIDE

01



HOW IT WORKS

A coenzyme central to ATP production and the substrate for sirtuins and PARPs; restoring it reactivates longevity pathways, improves mitochondrial efficiency, and lowers inflammation.

02



CELLULAR ENERGY

Increases mitochondrial energy production.


03



CLARITY & MOOD

Supports mental clarity, memory, and mood.

MOLECULAR STRUCTURE



MW $\approx$ 663.4 Da

1000 MG / 5 ML VIAL

Nicotinamide Adenine
Dinucleotide

04



PAIRS WELL WITH

MOTS-c ·
mitochondrial
Epitalon · longevity
Thymosin Alpha-1 ·
systemic repair


Complementary agents
commonly combined in
clinical protocols.

05



LONGEVITY

Promotes cellular
repair and may extend
healthspan via sirtuins.



MITOCHONDRIAL &
CELLULAR ENERGY



SAFETY NOTES

Well tolerated when used appropriately. Rapid infusion can cause transient flushing, chest tightness, nausea, or warmth; slower administration minimizes these. It is non-toxic and non-suppressive and suitable for long-term use.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

SS-3I

Stabilizing the engine of the cell.

CARDIOLIPIN-TARGETING TETRAPEPTIDE

01



HOW IT WORKS

Selectively binds cardiolipin in the inner mitochondrial membrane, protecting the cristae and electron transport chain; this improves ATP production and reduces reactive-oxygen-species leakage in stressed or aging cells.

02



CELLULAR ENERGY

Improves mitochondrial ATP production in stressed tissue.


03



OXIDATIVE STRESS

Reduces oxidative stress by limiting reactive oxygen species.

MOLECULAR STRUCTURE



C₃₂H₄₉N₉O₅

MW ≈ 639.8 Da

50 MG / 5 ML VIAL

Elamipretide
Cardiolipin-Targeting
Tetrapeptide

04



PAIRS WELL WITH

NAD⁺ · cellular energy
MOTS-c · mitochondrial signaling
Epitalon · longevity


Complementary agents commonly combined in clinical protocols.

05



ORGAN SUPPORT

Supports cardiac, renal, and neuronal function in mitochondrial dysfunction.



**MITOCHONDRIAL &
CELLULAR ENERGY**



SAFETY NOTES

Generally well tolerated in clinical study, with injection-site reactions (redness, irritation) the most common effect. Other reported effects are mild and may include headache or GI upset. It is non-suppressive and has shown a favorable profile across trials, though long-term human data are still developing. Use under clinician oversight in patients with significant cardiac or renal disease.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

Glutathione

The body's master antioxidant.

MASTER ANTIOXIDANT

01



HOW IT WORKS

Cycles between reduced and oxidized forms to neutralize reactive oxygen species and regenerate other antioxidants; in the liver it conjugates toxins for elimination and supports mitochondrial and immune function.

02



ANTIOXIDANT DEFENSE

Neutralizes free radicals and lowers oxidative stress.


03



DETOXIFICATION

Supports liver detoxification and clearance of toxins.

MOLECULAR STRUCTURE



C₁₀H₁₇N₃O₆S

MW ≈ 307.3 Da

**2000 MG / 10 ML ·
6000 MG / 30 ML VIAL**

Gamma-Glutamyl-Cysteinyl-
Glycine
Master Antioxidant

04



PAIRS WELL WITH

NAD+ · cellular energy
SS-31 · mitochondrial protection
Epitalon · antioxidant longevity


Complementary agents commonly combined in clinical protocols.

05



SKIN & RECOVERY

Promotes skin brightness and supports cellular recovery.



**MITOCHONDRIAL &
CELLULAR ENERGY**



SAFETY NOTES

Glutathione is well tolerated and naturally present in the body. Injectable use is generally safe, with occasional mild effects such as transient lightheadedness or injection-site irritation. Rare reports of skin lightening beyond the intended area exist with prolonged intravenous use. Patients with asthma should use caution, as inhaled forms have been associated with bronchospasm. Sterility and quality of the compounded preparation are essential.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

BPC-157

Repair, carried through every tissue.

BODY PROTECTION COMPOUND

01



HOW IT WORKS

Drives healing through several pathways at once: upregulates growth-hormone receptors in tendon fibroblasts and activates the VEGFR2-Akt-eNOS axis for angiogenesis and cell migration.

02



TISSUE REPAIR

Accelerates repair of tendon, ligament, muscle, skin, and bone in animal models.


03



GUT PROTECTION

Protects and stabilizes the gastrointestinal lining; supports fistula and colitis healing.

MOLECULAR STRUCTURE



$C_{62}H_{98}N_{16}O_{22}$

MW $\approx$ 1,419.6 Da

10 MG / 5 ML VIAL

Pentadecapeptide
Body Protection Compound

04



PAIRS WELL WITH

TB-500 · soft-tissue synergy
GHK-Cu · skin and repair
KPV · gut barrier


Complementary agents commonly combined in clinical protocols.

05



NEURAL RECOVERY

Counters neural injury and supports recovery after traumatic insult.



TISSUE REPAIR & RECOVERY



SAFETY NOTES

Extremely well tolerated across human and animal models, with no known toxicity at therapeutic levels and no suppression of natural healing. Infrequent effects include nausea, mild headache, transient dizziness, or injection-site irritation. Use caution in patients on anticoagulant or antihypertensive medications, as BPC-157 may influence vascular healing and blood flow. Clinical evidence in humans remains limited.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

TB-500

Mobility and repair, signaled system-wide.

THYMOSIN BETA-4 ACTIVE FRAGMENT

01



HOW IT WORKS

Regulates actin to upregulate cell motility, differentiation, and angiogenesis; raises VEGF and matrix metalloproteinases to build vessels and remodel tissue, and lowers inflammatory cytokines.

02



INJURY RECOVERY

Accelerates recovery from muscle, tendon, and ligament injury.


03



FLEXIBILITY

Improves flexibility and reduces connective-tissue stiffness.

MOLECULAR STRUCTURE



C₂₁₂H₃₅₀N₅₆O₇₈S

MW ≈ 4,963.5 Da

10 MG / 5 ML VIAL

Thymosin Beta-4 Active
Fragment

04



PAIRS WELL WITH

BPC-157 · soft-tissue synergy
GHK-Cu · skin and repair
IGF-1 LR3 · anabolic repair


Complementary agents commonly combined in clinical protocols.

05



VASCULAR REPAIR

Supports angiogenesis and cardiovascular repair.



TISSUE REPAIR & RECOVERY



SAFETY NOTES

Generally well tolerated. Mild effects may include fatigue, lightheadedness, or transient flu-like symptoms; injection-site reactions are rare. No known toxicity or carcinogenicity is associated with TB-500, and it does not suppress natural hormone production.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

GHK-Cu

The copper signal behind collagen, skin, and hair renewal.

COPPER TRIPEPTIDE

01



HOW IT WORKS

Delivers bioavailable copper and drives collagen, elastin, and glycosaminoglycan synthesis; at the genetic level it resets hundreds of repair-related genes and raises antioxidant enzymes.

02



SKIN & COLLAGEN

Improves elasticity, softens fine lines, and lightens hyperpigmentation.


03



WOUND HEALING

Accelerates wound, burn, and ulcer repair; supports post-surgical recovery.

MOLECULAR STRUCTURE



MW $\approx$ 403.9 Da

50 MG / 5 ML VIAL

Glycyl-Histidyl-Lysine
Copper Tripeptide

04



PAIRS WELL WITH

BPC-157 · tissue repair
TB-500 · connective tissue
Sermorelin · skin and collagen


Complementary agents commonly combined in clinical protocols.

05



HAIR & FOLLICLE

Stimulates follicular repair for hair regrowth and thickening.



TISSUE REPAIR & RECOVERY



SAFETY NOTES

Extremely well tolerated in both topical and injectable forms, with no major side effects documented in clinical or preclinical use. Mild injection-site irritation, headache, or fatigue can occur in sensitive patients. Because GHK-Cu binds copper, monitor copper levels during extended systemic use; patients with copper-metabolism disorders or active malignancy should use it only under clinician supervision.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

Thymosin Alpha-1

Recalibrating immune intelligence.

IMMUNOREGULATORY PEPTIDE

01



HOW IT WORKS

A biological response modifier that activates T-lymphocytes and natural killer cells and raises interleukin-2 and interferon-gamma, while calming pro-inflammatory cytokines to rebalance immunity.

02



IMMUNE DEFENSE

Strengthens immune defense during infection.


03



SURVEILLANCE

Enhances immune surveillance of abnormal cells.

MOLECULAR STRUCTURE



C₁₂₉H₂₁₅N₃₃O₅₅

MW ≈ 3,108.3 Da

10 MG / 5 ML VIAL

Immunoregulatory Peptide

04



PAIRS WELL WITH

LL-37 · antimicrobial
KPV · anti-inflammatory
BPC-157 · recovery

Complementary agents commonly combined in clinical protocols.

05



INFLAMMATION

Calms inflammation in autoimmune conditions.



**INFLAMMATION &
IMMUNE DEFENSE**



SAFETY NOTES

An excellent safety profile with few adverse effects even in long-term use. Mild effects may include fatigue, slight injection-site irritation, or transient headache. Acting as a biologic normalizer, it does not overstimulate the immune system and may be used preventively or acutely.



KPV

THE ACTIVE PHARM

EST. 2024

503A PHARMACY COMPOUNDED

Inflammation, quieted at the source.

ALPHA-MSH FRAGMENT

01



HOW IT WORKS

Inhibits activation of NF-κB, the transcription factor that triggers inflammatory cytokine production. The result is reduced excess inflammation with normal immune defense preserved, supporting healthy inflammatory responses throughout the body.


02



GUT BARRIER

Promotes gastrointestinal and gut-barrier health.

MOLECULAR STRUCTURE



C₁₇H₃₂N₆O₄

MW ≈ 384.5 Da

10 MG / 5 ML VIAL

Lysine-Proline-Valine
alpha-MSH Fragment

03



PAIRS WELL WITH

BPC-157 · gut lining and tissue repair
Thymosin Alpha-1 · immune modulation
LL-37 · antimicrobial support

Complementary agents commonly combined in clinical protocols.

04



SKIN & TISSUE

Calms skin inflammation and assists tissue recovery.



INFLAMMATION & IMMUNE DEFENSE



SAFETY NOTES

Generally well tolerated. Possible effects are usually mild: transient headache, fatigue, GI upset, or injection-site irritation in sensitive patients. Because it modulates inflammatory signaling, use caution in patients with highly reactive immune profiles and monitor for unexpected sensitivity.



THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

LL-37

Innate defense, broadly armed.

CATHELICIDIN-DERIVED ANTIMICROBIAL PEPTIDE

01



HOW IT WORKS

Its cationic, amphipathic structure disrupts microbial membranes for broad-spectrum killing, while recruiting immune cells, tuning cytokines, and promoting angiogenesis and re-epithelialization.

02



BROAD DEFENSE

Broad-spectrum action against bacteria, viruses, and fungi.


03



WOUND REPAIR

Supports wound healing and tissue regeneration.

MOLECULAR STRUCTURE



C₂₀₄H₃₄₅N₅₅O₄₆

MW ≈ 4,304.3 Da

5 MG / 5 ML VIAL

Cathelicidin-Derived
Antimicrobial Peptide

04



PAIRS WELL WITH

Thymosin Alpha-1 · immune modulation
KPV · anti-inflammatory
BPC-157 · recovery

Complementary agents commonly combined in clinical protocols.

05



SKIN & BIOFILM

Calms inflammatory skin conditions and disrupts biofilms.



**INFLAMMATION &
IMMUNE DEFENSE**



SAFETY NOTES

Typically well tolerated but can cause injection-site irritation, flushing, or mild inflammation. Because it stimulates immune activity, use caution in patients with autoimmune disorders or high baseline inflammation. Long-term use is not recommended without provider oversight.



Semax

Focus and neural resilience.

THE ACTIVE PHARM

EST. 2024

503A PHARMACY COMPOUNDED

ACTH(4-10) HEPTAPEPTIDE ANALOG

01



HOW IT WORKS

Inhibits enkephalin breakdown and enhances dopamine and serotonin transmission; raises BDNF, nerve growth factor, and VEGF for neuroplasticity and cerebral blood flow, without raising heart rate.

02



FOCUS & CLARITY

Enhances focus, learning, and mental clarity.


03



DRIVE

Increases dopamine signaling for drive and motivation.

MOLECULAR STRUCTURE



C₃₇H₅₁N₉O₁₀

MW ≈ 781.9 Da

10 MG / 5 ML VIAL

ACTH(4-10) Heptapeptide
Analog

04



PAIRS WELL WITH

Selank ·
complementary
nootropic
NAD+ · neural energy


Complementary agents
commonly combined in
clinical protocols.

05



NEUROPROTECTION

Provides
neuroprotection after
stress, injury, or fatigue.



**COGNITIVE FUNCTION &
NEUROPROTECTION**



SAFETY NOTES

Extremely well tolerated, with minimal reported side effects. Occasional users may notice mild euphoria, nasal irritation, or light sensitivity. It is non-addictive and does not disturb sleep or appetite when used appropriately. Administration is typically intranasal.



Selank

Calm without sedation.

THE ACTIVE PHARM

EST. 2024

503A PHARMACY COMPOUNDED

TUFTSIN-DERIVED HEPTAPEPTIDE

01



HOW IT WORKS

Modulates the GABAergic system, enhancing natural GABA sensitivity and release for calm without sedation; also raises serotonin and dopamine turnover and BDNF expression.

02



ANXIETY & MOOD

Reduces anxiety and tension and improves emotional resilience.


03



FOCUS

Enhances focus and mental clarity.

MOLECULAR STRUCTURE



C₃₃H₅₇N₁₁O₉

MW ≈ 751.9 Da

10 MG / 5 ML VIAL

Tuftsins-Derived
Heptapeptide

04



PAIRS WELL WITH

Semax ·
complementary
nootropic
DSIP · sleep and stress


Complementary agents
commonly combined in
clinical protocols.

05



CLEAN PROFILE

Non-sedating and non-
habit-forming.



**COGNITIVE FUNCTION &
NEUROPROTECTION**



SAFETY NOTES

A strong safety profile in animal and human studies, with no sedation, withdrawal, or abuse potential. Side effects are rare and may include mild nasal irritation or transient headache in sensitive users. It is typically administered intranasally.



Epitalon

Tuned to the clock of cellular aging.

THE ACTIVE PHARM

EST. 2024

503A PHARMACY COMPOUNDED

PINEAL TETRAPEPTIDE (AEDG)

01



HOW IT WORKS

Activates telomerase to help preserve telomere length, regulates gene expression in stem and neural cells, raises antioxidant enzymes, and helps restore melatonin and circadian rhythm.

02



CELLULAR LONGEVITY

Supports telomere maintenance and cellular longevity.


03



CIRCADIAN RHYTHM

Normalizes circadian rhythm and melatonin production.

MOLECULAR STRUCTURE



$C_{14}H_{22}N_4O_9$

MW $\approx$ 390.3 Da

10 MG / 5 ML VIAL

Pineal Tetrapeptide (AEDG)

04



PAIRS WELL WITH

NAD+ · longevity pathways
DSIP · sleep
Thymosin Alpha-1 · immune resilience


Complementary agents commonly combined in clinical protocols.

05



ANTIOXIDANT

Provides antioxidant and neuroprotective effects.



LONGEVITY, SLEEP &
CIRCADIAN



SAFETY NOTES

A strong safety profile in animal and early human studies, with minimal reported adverse effects. Although telomerase activation raises theoretical concerns about uncontrolled cell growth, available studies suggest Epitalon may suppress rather than promote certain cancers. Long-term use is best approached under clinical supervision.



DSIP

Deep sleep, gently restored.

DELTA SLEEP-INDUCING PEPTIDE

01



HOW IT WORKS

Enhances slow-wave (delta) sleep and promotes sleep onset without acting as a sedative. It modulates the hypothalamus to balance hormonal rhythms, helps normalize cortisol, and shows antioxidant and neuroprotective activity in the central nervous system.


02



DEEP SLEEP

Enhances deep, restorative sleep and reduces stress-related disturbance.

MOLECULAR STRUCTURE



C₃₅H₄₈N₁₀O₁₅

MW ≈ 848.8 Da

10 MG / 5 ML VIAL

Delta Sleep-Inducing Peptide

03



PAIRS WELL WITH

Epitalon · circadian rhythm

Selank · stress and calm

Oxytocin · relaxation

Complementary agents commonly combined in clinical protocols.

04



CIRCADIAN BALANCE

Supports hormonal balance and circadian regulation.



LONGEVITY, SLEEP & CIRCADIAN



SAFETY NOTES

Non-sedating and non-habit-forming. Rare side effects include vivid dreams, headache, or mild grogginess. It should be avoided in patients with epilepsy or severe psychiatric disorders and used only under provider oversight.



PT-141

THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

Desire, restored at its source in the brain.

MELANOCORTIN AGONIST

01



HOW IT WORKS

Binds melanocortin receptors MC3R and MC4R in the hypothalamus, increasing dopamine release and engaging arousal and reward circuits; the effect is central rather than vascular.

02



DESIRE

Increases libido and sexual desire in both men and women.


03



FUNCTION

Supports erectile function independent of blood-flow mechanics.

MOLECULAR STRUCTURE



C₅₀H₆₈N₁₄O₁₀

MW ≈ 1,025.2 Da

10 MG / 5 ML VIAL

Bremelanotide
Melanocortin Agonist

04



PAIRS WELL WITH

Oxytocin · arousal and bonding
Kisspeptin · reproductive hormones

Complementary agents commonly combined in clinical protocols.

05



WHERE PDE5 FAILS

Effective where PDE5 inhibitors are not, with no blood-pressure impact.



**SEXUAL & REPRODUCTIVE
HEALTH**



SAFETY NOTES

Generally well tolerated. Side effects can include facial flushing, headache, nausea, and a transient rise in blood pressure, usually mild; subcutaneous injection carries fewer systemic effects than nasal forms. Not recommended for patients with uncontrolled hypertension or heart disease, or those taking dopamine agonists without clinician oversight.



Oxytocin

The chemistry of bonding.

THE ACTIVE PHARM
EST. 2024

503A PHARMACY COMPOUNDED

NONAPEPTIDE NEUROHORMONE

01



HOW IT WORKS

Binds receptors across the brain and body, modulating dopamine and serotonin to enhance trust and calm, amplifying arousal, and lowering amygdala activity to blunt fear and anxiety.

02



BONDING

Improves emotional bonding and connection.


03



CALM

Reduces anxiety and stress by calming the nervous system.

MOLECULAR STRUCTURE



$C_{43}H_{66}N_{12}O_{12}S_2$

MW $\approx$ 1,007.2 Da

10 MG / 5 ML VIAL

Nonapeptide Neurohormone

04



PAIRS WELL WITH

PT-141 · desire and arousal

Kisspeptin · hormonal signaling

DSIP · relaxation


Complementary agents commonly combined in clinical protocols.

05



AROUSAL

Enhances arousal and intimacy.



SEXUAL & REPRODUCTIVE
HEALTH



SAFETY NOTES

Generally well tolerated when used appropriately. Possible effects include mild headache, flushing, lightheadedness, or temporary emotional sensitivity; higher amounts may rarely cause hypotension or nausea. It should be avoided in pregnancy unless medically indicated. Intermittent use is preferred to avoid receptor desensitization.



Kisspeptin

The upstream switch of reproductive hormones.

THE ACTIVE PHARM

EST. 2024

503A PHARMACY COMPOUNDED

KISS1 FRAGMENT

01



HOW IT WORKS

Activates hypothalamic receptors that trigger GnRH release, driving the pituitary to release LH and FSH, which act on the gonads to support testosterone production and reproductive balance while preserving the body's own feedback.


02



HORMONE SIGNALING

Supports healthy testosterone levels and stimulates natural hormone pathways.

MOLECULAR STRUCTURE



C₆₃H₈₃N₁₇O₁₄

MW ≈ 1,302.5 Da

10 MG / 5 ML VIAL

Kisspeptin-10
KISS1 Fragment

03



PAIRS WELL WITH

PT-141 · desire and arousal
Oxytocin · intimacy

Complementary agents commonly combined in clinical protocols.

04



REPRODUCTIVE FUNCTION

Promotes reproductive function and libido.



SEXUAL & REPRODUCTIVE HEALTH



SAFETY NOTES

Generally well tolerated. Possible side effects include headache, flushing, nausea, or mild fatigue. Because it directly influences reproductive hormone signaling, patients with endocrine disorders or hormone-sensitive conditions should use it under clinician guidance.

Disclaimer & Professional Use

FDA Disclosure

These statements have not been evaluated by the Food and Drug Administration. These formulations are not intended to diagnose, treat, cure, or prevent any disease. Results may vary.

Sourcing & Compounding

All peptides and formulations offered by The Active Pharm are compounded exclusively through 503A pharmacies operating under licensed pharmacist oversight, in accordance with applicable state and federal compounding standards.

General Reference

This edition is provided for general educational reference and does not include dosing, titration, or cycling guidance. All dosing and administration decisions must come from the patient's licensed medical provider.

Pairing References

The "Pairs Well With" references describe agents commonly combined in practice and are not treatment recommendations.

Professional Use Notice

By placing an order, the purchaser affirms the following:

Licensure & Eligibility. The purchaser is a licensed medical provider, an authorized medical facility, or an entity operating under a provider's medical license.

Professional Use Only. All products are intended for professional or clinical use. They are made available only to licensed medical providers operating under direct medical oversight.

Responsibility & Oversight. The purchaser accepts full responsibility for the appropriate handling, storage, administration, and use of all products in accordance with their medical licensure, local regulations, and internal protocols. The Active Pharm does not provide diagnosis, treatment recommendations, or clinical guidance. All usage decisions remain solely at the discretion of the treating provider.

COMPOUNDED · 503A PHARMACY NETWORK · CLINICIAN USE ONLY